

About ELZONRIS® (tagraxofusp-erzs)

ELZONRIS is a prescription medicine used to treat blastic plasmacytoid dendritic cell neoplasm (BPDCN) in adults and pediatric patients 2 years and older.

IMPORTANT SAFETY INFORMATION

ELZONRIS can cause serious side effects, including:

- **Capillary Leak Syndrome (CLS).** ELZONRIS can cause fluid to leak from small blood vessels into your body’s tissues. This is called “Capillary Leak Syndrome.” CLS can quickly cause you to have symptoms that may become life-threatening or fatal (ie, lead to death). **Get emergency medical help immediately if you develop any of the following symptoms:**
 - fast weight gain
 - swelling of your face, arms, hands, legs, or feet
 - shortness of breath or difficulty breathing
 - low blood pressure (dizziness or lightheadedness, headache, feeling tired, or shortness of breath)

Weigh yourself daily. Your healthcare provider will also check your weight and test your blood before you receive each dose of ELZONRIS and as needed during treatment.

- **Hypersensitivity reactions** may occur with ELZONRIS. Symptoms may include rash, itching (pruritus), wheezing, or swelling in your face, including around your eyes and/or in and around your mouth
- **Liver damage** is usually detected through blood tests. Symptoms may include feeling tired (fatigue), loss of appetite, yellowing of your skin or the whites of your eyes (jaundice), or upper right abdominal pain or discomfort

Your healthcare provider will periodically test your blood while you are on ELZONRIS to check for liver damage. Contact your healthcare provider immediately if you have any of these symptoms.

Getting medical treatment right away may help keep these problems from becoming more serious.

If you have any side effects during treatment with ELZONRIS, your healthcare provider may hold your treatment for a period of time or completely stop your treatment with ELZONRIS.

The most common side effects of ELZONRIS include CLS, nausea, feeling tired (fatigue), fever, swelling in your legs or feet, and weight gain.

These are not all of the possible side effects of ELZONRIS. If any new side effects start or an existing one gets worse, contact your healthcare provider immediately. For more information, talk to your treatment team.

Be sure to tell your treatment team about:

- all of your medical conditions, including if you
 - are pregnant or plan to become pregnant. ELZONRIS may harm your unborn baby
 - If you are a female who can become pregnant, you should use effective birth control during ELZONRIS treatment and for 1 week after the last dose
 - Tell your healthcare provider right away if you become pregnant during treatment with ELZONRIS
 - are breastfeeding or plan to breastfeed. It is not known if ELZONRIS passes into breast milk. You and your healthcare provider should decide if you will receive ELZONRIS or breastfeed. You should not do both
- all of the medicines you take, including prescription medicines, over-the-counter medicines, vitamins, and herbal supplements

You can report any side effects to Stemline Therapeutics, Inc. at 1-877-332-7961 or contact the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see Full Prescribing Information, including **Boxed WARNING**, for ELZONRIS to learn more.

The risk information provided here is not comprehensive. To learn more, talk about ELZONRIS (tagraxofusp-erzs) with your healthcare provider or pharmacist. The FDA-approved product labeling can be found accompanying this booklet.



How Stemline ARC® can help



Stemline ARC offers ongoing support from ARC Patient Advocates during your treatment. Talk to your healthcare team about enrolling in Stemline ARC.

ARC Patient Advocates provide a single point of contact.

ARC Patient Advocates provide an overview of support services, confirm ELZONRIS coverage and financial assistance, share helpful resources, and answer questions throughout your treatment journey.

This supplemental support is not intended to replace discussions between you and your healthcare provider.

Get more information about Stemline ARC

- Talk to your healthcare team about enrolling
- Call 1-833-4-STEMLINE (1-833-478-3654) from 9:00 AM to 6:00 PM EST, Monday through Friday

Please see Important Safety Information, including **Boxed WARNING**, on page 11 and accompanying Full Prescribing Information, including **Boxed WARNING**, for ELZONRIS to learn more.

ELZONRIS is a registered trademark of Stemline Therapeutics, Inc.
© 2025 Stemline Therapeutics, Inc., a Menarini Group Company.
All rights reserved. 05/25 MAT-US-TAG-00246



My ELZONRIS Treatment Tracker

About ELZONRIS® (tagraxofusp-erzs)

ELZONRIS is a prescription medicine used to treat blastic plasmacytoid dendritic cell neoplasm (BPDCN) in adults and pediatric patients 2 years and older.

ELZONRIS can cause serious side effects, including:

- **Capillary Leak Syndrome (CLS).** ELZONRIS can cause fluid to leak from small blood vessels into your body’s tissues. This is called “Capillary Leak Syndrome.” CLS can quickly cause you to have symptoms that may become life-threatening or fatal (ie, lead to death).

Please see Important Safety Information, including **Boxed WARNING**, on page 11 and accompanying Full Prescribing Information, including **Boxed WARNING**, for ELZONRIS to learn more.



What you may experience while receiving ELZONRIS



When you are treated with ELZONRIS for your blastic plasmacytoid dendritic cell neoplasm (BPDCN), you may experience side effects from the treatment, or symptoms from your cancer or other health conditions. Be sure to speak with your healthcare team about how you feel while taking ELZONRIS, as they may be able to help.

The most common side effects that may occur with ELZONRIS include **capillary leak syndrome (CLS), nausea, fatigue** (feeling tired), **fever, edema** (when fluid builds up in the body), **and weight increase**. These are not the only side effects that may occur. You can get more information from your healthcare team.



Call your healthcare provider about any symptoms that concern you, or with any questions about your treatment with ELZONRIS.

PI POCKET

PI POCKET



ELZONRIS may cause a serious side effect known as **CLS**. Call your treatment team immediately if you have any signs or symptoms of CLS, including:

- fast weight gain
- swelling of your face, arms, hands, legs, or feet
- shortness of breath or difficulty breathing
- low blood pressure (dizziness or lightheadedness, headache, feeling tired, or shortness of breath)

Other common side effects




Common side effects	How you may feel
Nausea	<ul style="list-style-type: none">• You may feel sick to your stomach or as if you need to throw up• You may have no appetite
Fatigue	<ul style="list-style-type: none">• You may feel tired or sleepy, even after resting
Fever	<ul style="list-style-type: none">• You may find yourself sweating or feeling cold for no reason• You may have a headache or feel achy all over your body• You may feel weak or dehydrated
Edema	<ul style="list-style-type: none">• Your face, arms, hands, legs, or feet may feel swollen or puffy
Weight increase	<ul style="list-style-type: none">• You may gain an unusual amount of weight in a short period, while not necessarily eating more than usual

Note: These are not the only side effects that can occur. Talk to your healthcare team about any other side effects that you may be experiencing after starting treatment with ELZONRIS.

How to use the ELZONRIS Treatment Tracker


Use this tracker if you have any side effects while you receive treatment with ELZONRIS.

1




Fill in the date at the top of the column.

2




Note any symptoms you may be feeling in the column on the left. Describe each symptom with as many details as possible.

3



Use the scale to describe how a side effect feels, from 1 (mild) to 5 (severe).

4



Bring this tracker to your next doctor's visit and share the information.

Here is an example of how to fill out your symptom tracker.

ELZONRIS Treatment Tracker							
Date ▶	Feb. 12	Feb. 14	Feb. 15	Feb. 17			
Felt tired	1			1			
Felt sick to my stomach		2	1	3			

ELZONRIS Treatment Tracker

Date ▶							



If any side effect gets worse from one day to the next, it is very important to call your doctor immediately.

ELZONRIS Treatment Tracker

Date ▶							



If any side effect gets worse from one day to the next,
it is very important to call your doctor immediately.

ELZONRIS Treatment Tracker

Date ▶							



If any side effect gets worse from one day to the next,
it is very important to call your doctor immediately.

ELZONRIS Treatment Tracker

Date ▶							



If any side effect gets worse from one day to the next,
it is very important to call your doctor immediately.

ELZONRIS Treatment Tracker

Date ▶							



If any side effect gets worse from one day to the next,
it is very important to call your doctor immediately.

ELZONRIS Treatment Tracker

Date ▶							



If any side effect gets worse from one day to the next,
it is very important to call your doctor immediately.

Please see Important Safety Information, **including Boxed WARNING**, on page 11 and accompanying Full Prescribing Information, **including Boxed WARNING**, for ELZONRIS to learn more.